

This material was developed by JAMP Pharma, as part of the risk minimization plan for JAMP Teriflunomide. This material is not intended for promotional use.

Patient's name:

.....  
Date JAMP Teriflunomide first prescribed: \_\_/\_\_/\_\_

Hospital/Center name:

.....  
Treating doctor's/HCP name:

.....  
Treating doctor's/HCP phone number:

 For further information, please contact JAMP Medical Information Services at 1-866-399-9091 or email: [medinfo@jamppharma.com](mailto:medinfo@jamppharma.com). Website: [www.jampinfo.com](http://www.jampinfo.com) 

V2.0 Date of Preparation: March 2022.

---

## PATIENT CARD

This patient card provides important information on the risks of JAMP Teriflunomide (teriflunomide). Please show this card to any doctor or healthcare professional (HCP) involved in your medical care (e.g. in case of an emergency). You should also read the Patient Medication Information section of the Product Monograph for further information.

If you notice any side effects talk to your doctor, pharmacist or nurse. You can also report side effects directly, to Health Canada: via [canada.ca/medeffect](http://canada.ca/medeffect); Tel: 1-866-234-2345.

Or contacting us directly: JAMP Medical Information Services. Call us at 1-866-399-9091 or email [medinfo@jamppharma.com](mailto:medinfo@jamppharma.com). Website: [www.jampinfo.com](http://www.jampinfo.com)

### **Important side effects**

Teriflunomide reduces the activity of the immune system (immunomodulator). In some people, teriflunomide can cause liver damage (hepatitis) and it may also reduce the production of white blood cells that fight infection (neutrophils) and platelets that are involved in blood clotting. Your liver function tests, and blood pressure should be checked regularly during teriflunomide treatment, and your full blood count should be checked if necessary. These tests should also be checked before starting treatment.

### **If you have any of the following side effects, please contact your doctor/HCP immediately:**

- Yellow skin or yellowing of the whites of your eyes (jaundice), unexplained nausea or vomiting, abdominal pain or darker urine than normal. These are the symptoms of a liver problem.
- Signs of an infection including, pain on passing urine, confusion, high temperature (fever), cough, swollen glands.

### **For women of childbearing potential including their caregivers**

---

- JAMP Teriflunomide should not be used in pregnancy or in women of child-bearing potential if they are not using effective contraception because it can cause serious birth defects.
- Do not start JAMP Teriflunomide when you are pregnant, or you think you may be pregnant. Your doctor/HCP should administer a pregnancy test before you start taking JAMP Teriflunomide to make sure.
- Effective contraception should be used during and after JAMP Teriflunomide treatment until the blood levels are low. Your doctor/HCP will provide counselling on the potential risks to an unborn baby and on the need for effective contraception.
- Tell your doctor/HCP if you want to change your method of contraception or plan to become pregnant after stopping treatment with JAMP Teriflunomide. You should also discuss with your doctor/HCP if you plan to or are breastfeeding.
- Contact your doctor/HCP immediately if you are pregnant, are late starting your period, or suspect you may be pregnant while taking JAMP Teriflunomide or within 2 years of stopping it. If the pregnancy is confirmed, your doctor/HCP may suggest treatment with certain medicines to speed up the removal of teriflunomide from your body, as this may decrease the risk to your baby. Your doctor/HCP will encourage enrolment in the JAMP Teriflunomide Enhanced Pharmacovigilance Pregnancy Active Surveillance Program conducted by JAMP Pharma: Phone: 1 866 399 9091. Email: [medinfo@jamppharma.com](mailto:medinfo@jamppharma.com) ; Program Website: [www.jampinfo.com](http://www.jampinfo.com)